

Patient: **SAMPLE**  
**PATIENT**

DOB:

Sex:

MRN:

**3301 Organix - Urine**

**Results Overview**

organic acids



**MITOCHONDRIAL  
DYSFUNCTION**



**TOXIC  
EXPOSURE**



**METHYLATION  
IMBALANCE**

**Functional Imbalance Scores**

Key

**0-4** : Minimal Need for Support

**5-7** : Moderate Need for Support

**8-10** : High Need for Support

Need for Mitochondrial Support	
<b>Mitochondrial Dysfunction</b>	
<b>8</b>	
FIGLU	●
Methylmalonic Acid	▲
Glutaric Acid	▲
Lactic Acid	●
Pyruvic Acid	●
Citric Acid	▲
cis-Aconitic Acid	▲
Isocitric Acid	▲
α-Ketoglutaric Acid	▲
Succinic Acid	▲
Malic Acid	▲
Adipic Acid	●
Suberic Acid	▲

Need for Reduced Exposure	
<b>Toxic Exposure</b>	
<b>8</b>	
α-Hydroxyisobutyric Acid	▲
α-Ketophenylacetic Acid	▲
Pyroglutamic Acid	▲
Orotic Acid	▲
Citric Acid	▲
cis-Aconitic Acid	▲
Isocitric Acid	▲
Glutaric Acid	▲

Need for Methylation Support	
<b>Methylation Imbalance</b>	
<b>7</b>	
Methylmalonic Acid	▲
FIGLU	●
Vanilmandelic Acid	▲
Creatinine	●



## Nutrient Need Overview

	Nutrient Need										DRI	Suggested Recommendations	Provider Recommendations
	0	1	2	3	4	5	6	7	8	9			
<b>Antioxidants</b>													
Glutathione													
<b>B-Vitamins</b>													
Thiamin - B1											1.1 mg	25 mg	
Riboflavin - B2											1.1 mg	50 mg	
Niacin - B3											14 mg	50 mg	
Pyridoxine - B6											1.3 mg	25 mg	
Biotin - B7											30 mcg	400 mcg	
Folate - B9											400 mcg	800 mcg	
Cobalamin - B12											2.4 mcg	1,000 mcg	
<b>Minerals</b>													
Magnesium											320 mg	800 mg	
Manganese											1.8 mg	5.0 mg	
Zinc											8 mg	10 mg	
<b>GI Support</b>													
Digestive Support/Enzymes												10,000 IU	
Microbiome Support/Probiotics												50 billion CFU	

Recommendations for age and gender-specific supplementation are set by comparing levels of nutrient functional need to optimal levels as described in the peer-reviewed literature. They are provided as guidance for short-term support of nutritional deficiencies only.

The Nutrient Need Overview is provided at the request of the ordering practitioner. Any application of it as a therapeutic intervention is to be determined by the ordering practitioner.



## Interpretation At-A-Glance

### Antioxidant Needs

#### Glutathione



- Glutathione (GSH) is composed of cysteine, glutamine & glycine. GSH is a source of sulfate and plays a key role in antioxidant activity and detoxification of toxins.
- GSH requirement is increased with high-fat diets, cigarette smoke, cystinuria, chronic alcoholism, chronic acetaminophen use, infection, inflammation and toxic exposure.
- Deficiency may result in oxidative stress & damage, impaired detoxification, altered immunity, macular degeneration and increased risk of chronic illness.
- Food sources of GSH precursors include meats, poultry, fish, soy, corn, nuts, seeds, wheat germ, milk and cheese.

#### KEY

- Function of Nutrient
- Cause of Deficiency
- Complications of Deficiency
- Food Sources of Nutrient

## Interpretation At-A-Glance

### B-Vitamin Needs

#### Thiamin - B1



5

- B1 is a required cofactor for enzymes involved in energy production from food, and for the synthesis of ATP, GTP, DNA, RNA and NADPH.
- Low B1 can result from chronic alcoholism, diuretics, digoxin, oral contraceptives and HRT, or large amounts of tea & coffee (contain anti-B1 factors).
- B1 deficiency may lead to dry beriberi (e.g., neuropathy, muscle weakness), wet beriberi (e.g., cardiac problems, edema), encephalopathy or dementia.
- Food sources include lentils, whole grains, wheat germ, Brazil nuts, peas, organ meats, brewer's yeast, blackstrap molasses, spinach, milk & eggs.

#### Riboflavin - B2



10

- B2 is a key component of enzymes involved in antioxidant function, energy production, detoxification, methionine metabolism and vitamin activation.
- Low B2 may result from chronic alcoholism, some anti-psychotic medications, oral contraceptives, tricyclic antidepressants, quinacrine or adriamycin.
- B2 deficiency may result in oxidative stress, mitochondrial dysfunction, low uric acid, low B3 or B6, high homocysteine, anemia or oral & throat inflammation.
- Food sources include milk, cheese, eggs, whole grains, beef, chicken, wheat germ, fish, broccoli, asparagus, spinach, mushrooms and almonds.

#### Niacin - B3



8

- B3 is used to form NAD and NADP, involved in energy production from food, fatty acid & cholesterol synthesis, cell signaling, DNA repair & cell differentiation.
- Low B3 may result from deficiencies of tryptophan (B3 precursor), B6, B2 or Fe (cofactors in B3 production), or from long-term isoniazid or oral contraceptive use.
- B3 deficiency may result in pellagra (dermatitis, diarrhea, dementia), neurologic symptoms (e.g., depression, memory loss), bright red tongue or fatigue.
- Food sources include poultry, beef, organ meats, fish, whole grains, peanuts, seeds, lentils, brewer's yeast and lima beans.

#### Pyridoxine - B6



6

- B6 (as P5P) is a cofactor for enzymes involved in glycogenolysis & gluconeogenesis, and synthesis of neurotransmitters, heme, B3, RBCs and nucleic acids.
- Low B6 may result from chronic alcoholism, long-term diuretics, estrogens (oral contraceptives and HRT), anti-TB meds, penicillamine, L-DOPA or digoxin.
- B6 deficiency may result in neurologic symptoms (e.g., irritability, depression, seizures), oral inflammation, impaired immunity or increased homocysteine.
- Food sources include poultry, beef, beef liver, fish, whole grains, wheat germ, soybean, lentils, nuts & seeds, potato, spinach and carrots.

#### Biotin - B7



8

- Biotin is a cofactor for enzymes involved in functions such as fatty acid synthesis, mitochondrial FA oxidation, gluconeogenesis and DNA replication & transcription.
- Deficiency may result from certain inborn errors, chronic intake of raw egg whites, long-term TPN, anticonvulsants, high-dose B5, sulfa drugs & other antibiotics.
- Low levels may result in neurologic symptoms (e.g., paresthesias, depression), hair loss, scaly rash on face or genitals or impaired immunity.
- Food sources include yeast, whole grains, wheat germ, eggs, cheese, liver, meats, fish, wheat, nuts & seeds, avocado, raspberries, sweet potato and cauliflower.

#### Folate - B9



6

- Folate plays a key role in coenzymes involved in DNA and SAMe synthesis, methylation, nucleic acids & amino acid metabolism and RBC production.
- Low folate may result from alcoholism, high-dose NSAIDs, diabetic meds, H2 blockers, some diuretics and anti-convulsants, SSRIs, methotrexate, trimethoprim, pyrimethamine, triamterene, sulfasalazine or cholestyramine.
- Folate deficiency can result in anemia, fatigue, low methionine, increased homocysteine, impaired immunity, heart disease, birth defects and CA risk.
- Food sources include fortified grains, green vegetables, beans & legumes.

#### Cobalamin - B12



8

- B12 plays important roles in energy production from fats & proteins, methylation, synthesis of hemoglobin & RBCs, and maintenance of nerve cells, DNA & RNA.
- Low B12 may result from alcoholism, malabsorption, hypochlorhydria (e.g., from atrophic gastritis, H. pylori infection, pernicious anemia, H2 blockers, PPIs), vegan diets, diabetic meds, cholestyramine, chloramphenicol, neomycin or colchicine.
- B12 deficiency can lead to anemia, fatigue, neurologic symptoms (e.g., paresthesias, memory loss, depression, dementia), methylation defects or chromosome breaks.
- Food sources include shellfish, red meat, poultry, fish, eggs, milk and cheese.

### KEY

- Function of Nutrient
- Cause of Deficiency
- Complications of Deficiency
- Food Sources of Nutrient



## Interpretation At-A-Glance

### Mineral Needs

#### Magnesium

8

- Magnesium is involved in >300 metabolic reactions. Key areas include energy production, bone & ATP formation, muscle & nerve conduction and cell signaling.
- Deficiency may occur with malabsorption, alcoholism, hyperparathyroidism, renal disorders (wasting), diabetes, diuretics, digoxin or high doses of zinc.
- Low Mg may result in muscle weakness/spasm, constipation, depression, hypertension, arrhythmias, hypocalcemia, hypokalemia or personality changes.
- Food sources include dark leafy greens, oatmeal, buckwheat, unpolished grains, chocolate, milk, nuts & seeds, lima beans and molasses.

#### Manganese

6

- Manganese plays an important role in antioxidant function, gluconeogenesis, the urea cycle, cartilage & bone formation, energy production and digestion.
- Impaired absorption of Mn may occur with excess intake of Fe, Ca, Cu, folic acid, or phosphorous compounds, or use of long-term TPN, Mg-containing antacids or laxatives.
- Deficiency may result in impaired bone/connective tissue growth, glucose & lipid dysregulation, infertility, oxidative stress, inflammation or hyperammonemia.
- Food sources include whole grains, legumes, dried fruits, nuts, dark green leafy vegetables, liver, kidney and tea.

#### Zinc

0

- Zinc plays a vital role in immunity, protein metabolism, heme synthesis, growth & development, reproduction, digestion and antioxidant function.
- Low levels may occur with malabsorption, alcoholism, chronic diarrhea, diabetes, excess Cu or Fe, diuretics, ACE inhibitors, H2 blockers or digoxin.
- Deficiency can result in hair loss and skin rashes, also impairments in growth & healing, immunity, sexual function, taste & smell and digestion.
- Food sources include oysters, organ meats, soybean, wheat germ, seeds, nuts, red meat, chicken, herring, milk, yeast, leafy and root vegetables.

### KEY

- Function of Nutrient
- Cause of Deficiency
- Complications of Deficiency
- Food Sources of Nutrient



## Interpretation At-A-Glance

### Microbiome & Digestive Support

#### Microbiome Support/Probiotics



8

- Probiotics have many functions. These include: production of some B vitamins and vitamin K; enhance digestion & absorption; decrease severity of diarrheal illness; modulate of immune function & intestinal permeability.
- Alterations of gastrointestinal microflora may result from C-section delivery, antibiotic use, improved sanitation, decreased consumption of fermented foods and use of certain drugs.
- Some of the diseases associated with microflora imbalances include: IBS, IBD, fibromyalgia, chronic fatigue syndrome, obesity, atopic illness, colic and cancer.
- Food sources rich in probiotics are yogurt, kefir and fermented foods.

#### Digestive Support/Enzymes



8

- Pancreatic enzymes are secreted by the exocrine glands of the pancreas and include protease/peptidase, lipase and amylase.
- Pancreatic exocrine insufficiency may be primary or secondary in nature. Any indication of insufficiency warrants further evaluation for underlying cause (i.e., celiac disease, small intestine villous atrophy, small bowel bacterial overgrowth).
- A high functional need for digestive enzymes suggests that there is an impairment related to digestive capacity.
- Determining the strength of the pancreatic enzyme support depends on the degree of functional impairment. Supplement potency is based on the lipase units present in both prescriptive and non-prescriptive agents.

### Functional Imbalances

#### Mitochondrial Dysfunction



8

- Mitochondria are a primary site of generation of reactive oxygen species. Oxidative damage is considered an important factor in decline of physiologic function that occurs with aging and stress.
- Mitochondrial defects have been identified in cardiovascular disease, fatigue syndromes, neurologic disorders such as Parkinson's and Alzheimer's disease, as well as a variety of genetic conditions. Common nutritional deficiencies can impair mitochondrial efficiency.

#### Need for Methylation



7

- Methylation is an enzymatic process that is critical for both synthesis and inactivation. DNA, estrogen and neurotransmitter metabolism are all dependent on appropriate methylation activity.
- B vitamins and other nutrients (methionine, magnesium, selenium) functionally support catechol-O-methyltransferase (COMT), the enzyme responsible for methylation.

#### Toxic Exposure



8


- Methyl tert-Butyl Ether (MTBE) is a common gasoline additive used to increase octane ratings, and has been found to contaminate ground water supplies where gasoline is stored. Inhalation of MTBE may cause nose and throat irritation, as well as headaches, nausea, dizziness and mental confusion. Animal studies suggest that drinking MTBE may cause gastrointestinal irritation, liver and kidney damage and nervous system effects.
- Styrene is classified by the US EPA as a "potential human carcinogen," and is found widely distributed in commercial products such as rubber, plastic, insulation, fiberglass, pipes, food containers and carpet backing.
- Levels of these toxic substances should be examined within the context of the body's functional capacity for methylation and need for glutathione.

### KEY

- Function of Nutrient
- Cause of Deficiency
- Complications of Deficiency
- Food Sources of Nutrient



## Oxidative Stress & Mitochondrial Dysfunction



All biomarkers reported in mmol/mol creatinine unless otherwise noted.



Organic Acids			
Malabsorption & Dysbiosis Markers		Vitamin Markers	
<b>Malabsorption Markers</b>		<b>Branched-Chain Catabolites (B1, B2, B3, ALA)</b>	
	Reference Range		Reference Range
Indoleacetic Acid	40.0	α-Ketoadipic Acid	1.9
Phenylacetic Acid	<dl	α-Ketoisovaleric Acid	<dl
<b>Dysbiosis Markers</b>		<b>Methylation Markers (Folate, B12)</b>	
Dihydroxyphenylpropionic Acid (DHPPA)	18.9	Formiminoglutamic Acid (FIGlu)	0.6
3-Hydroxyphenylacetic Acid	5.8	Methylmalonic Acid	2.7
4-Hydroxyphenylacetic Acid	19	<b>Biotin Markers</b>	
Benzoic Acid	0.10	3-Hydroxypropionic Acid	15
Hippuric Acid	340	3-Hydroxyisovaleric Acid	25
<b>Yeast / Fungal Dysbiosis Markers</b>		<b>Neurotransmitter Metabolites</b>	
D-Arabinitol	14	<b>Kynurenine Markers (Vitamin B6)</b>	
Citramalic Acid	2.6	Kynurenic Acid	8.7
Tartaric Acid	5	Quinolinic Acid	4.2
<b>Cellular Energy &amp; Mitochondrial Markers</b>		<b>Catecholamine Markers</b>	
<b>Fatty Acid Metabolism</b>		<b>Homovanillic Acid</b>	
Adipic Acid	1.3	Vanilmandelic Acid	4.2
Suberic Acid	2.5	3-Methyl-4-OH-phenylglycol	0.13
<b>Carbohydrate Metabolism</b>		<b>Serotonin Markers</b>	
Pyruvic Acid	21	5-OH-indoleacetic Acid	24.2
Lactic Acid	10.8	<b>Toxin &amp; Detoxification Markers</b>	
α-Hydroxybutyric Acid	0.70	Pyroglutamic Acid	68
β-OH-Butyric Acid	1.4	α-Ketophenylacetic Acid (from Styrene)	0.33
β-OH-β-Methylglutaric Acid	9	α-Hydroxyisobutyric Acid (from MTBE)	7.5
<b>Energy Metabolism</b>		Orotic Acid	0.81
Citric Acid	760	Organic Acid Reference Ranges are Age Specific	
cis-Aconitic Acid	50		
Isocitric Acid	100		
α-Ketoglutaric Acid	100		
Succinic Acid	3.7		
Malic Acid	3.2		

Methodology: GCMS, LC/MS/MS, Alkaline Picrate, Colorimetric



Methodology: Colorimetric, thiobarbituric acid reactive substances (TBARS), Alkaline Picrate, Hexokinase/G-6-PDH, HPLC, GC/MS

Organic Acids			
Oxalate Markers	Reference Range	Creatinine Concentration	Reference Range
Glyceric Acid	23.7 3.5-16.4	1.8	3.1-19.5 mmol/L
Glycolic Acid	59 <= 67		
Oxalic Acid	66 <= 78		


All biomarkers reported in mmol/mol creatinine.

Oxidative Stress Markers	
Oxidative Damage	Reference Range
8-OHdG (urine)	<DL <= 15 mcg/g Creat.

The Oxidative Stress reference ranges are based on an adult population.

### Pathways

#### Methylation Markers



#### Branch-Chain Amino Acid Metabolism

